

## 兔抗 F7 多克隆抗体

中文名称: 兔抗 F7 多克隆抗体

英文名称: Anti-F7 rabbit polyclonal antibody

别 名: SPCA

储 存: 冷冻(-20℃)

抗 原: F7

宿 主: Rabbit

相关类别: 一抗

反应种属: Human

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

Xa, factor XIIa, or thrombin by minor proteolysis. Upon activation of the factor VII, a heavy chain containing a catalytic domain and a light chain containing 2 EGF-like domains are generated, and two chains are held togeth er by a disulfide bond. In the presence of factor III and

calcium ions, the activated factor then further activates the coagulation cascade by converting factor IX to fact or IXa and/or factor X to factor Xa. Defects in this gen

This gene encodes coagulation factor VII which is a vita min K-dependent factor essential for hemostasis. This fa ctor circulates in the blood in a zymogen form, and is converted to an active form by either factor IXa, factor

e can cause coagulopathy. Alternatively spliced transcrip



	t variants encoding multiple isoforms have been observ ed for this gene.
Applications:	ELISA, IHC
Name of antibody:	F7
Immunogen:	Fusion protein of human F7
Full name:	coagulation factor VII (serum prothrombin conversion a ccelerator)
Synonyms :	SPCA
SwissProt:	P08709
ELISA Recommended dilution:	1000-5000
IHC positive control:	Human thyroid cancer and human lung cancer
IHC Recommend dilution:	15-50

